

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
13 January 2005 (13.01.2005)

PCT

(10) International Publication Number
WO 2005/003116 A1

(51) International Patent Classification⁷: **C07D 333/56**

(21) International Application Number:
PCT/EP2004/051263

(22) International Filing Date: 28 June 2004 (28.06.2004)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
MI2003A001333 30 June 2003 (30.06.2003) IT

(71) Applicant (for all designated States except US): **ER-REGIERRE S.p.A.** [IT/IT]; Via Francesco Baracca 19, I-24060 SAN PAOLO D'ARGON (IT).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **FERRARI, Massimo** [IT/IT]; Via Antonio Locatelli 7, I-24069 CENATE SOTTO (IT). **ZINETTI, Fabrizio** [IT/IT]; Via Valle 7, I-24060 CASAZZA (IT). **BELOTTI, Paolo** [IT/IT]; Via Cesare Battisti 5, I-24060 SAN PAOLO D'ARGON (IT).

(74) Agent: **GERVASI, Gemma**; **NOTARBARTOLO & GERVASI S.p.A.**, Corso di Porta Vittoria 9, I-20122 MILAN (IT).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declaration under Rule 4.17:

— of inventorship (Rule 4.17(iv)) for US only

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: **PROCESS FOR PREPARING RALOXIFENE HYDROCHLORIDE**

(57) Abstract: Process for preparing raloxifene hydrochloride with a purity greater than 98% and low aluminium content comprising the following stages a) demethylation of 6-methoxy-2-(4-methoxyphenyl)benzo[b]thiophene in pyridine and hydrochloric acid to obtain 6-hydroxy-2-(4-hydroxyphenyl)benzo[b]thiophene in pyridine hydrochloride, b) acetylation of 6-hydroxy-2-(4-hydroxyphenyl)benzo[b]thiophene with an acetylating agent to obtain the corresponding 6-acetoxy-2-(4-acetoxyphenyl)benzo[b]thiophene, c) acylation of 6-acetoxy-2-(4-acetoxyphenyl)benzo[b]thiophene with 4-(2-piperidinoethoxy)benzoylchloride hydrochloride with aluminium trichloride in halogenated solvent to obtain 6-acetoxy-2-(4-acetoxyphenyl)-3-[4-(2-piperidinoethoxy)benzoyl]-benzo[b]thiophene, d) hydrolysis of 6-acetoxy-2-(4-acetoxyphenyl)-3-[4-(2-piperidinoethoxy)benzoyl]benzo[b]thiophene according to the following operating conditions: d1) treatment of 6-acetoxy-2-(4-acetoxyphenyl)-3-[4-(2-piperidinoethoxy)benzoyl]benzo[b]thiophene with alkaline hydroxide in alcohol solvent, d2) acidification of the product obtained in the preceding stage (d1) with a strong acid, to obtain the corresponding raloxifene salt with the strong acid, characterised in that the strong acid used in stage (d2) is concentrated hydrochloric acid.

WO 2005/003116 A1


PROCESS FOR PREPARING RALOXIFENE HYDROCHLORIDE

Field of the invention

The present invention relates to a process for preparing raloxifene and in particular high purity raloxifene hydrochloride with high yields.

State of the art


Raloxifene and in particular the relating hydrochloride salt, characterised by the following formula (I):



15 is an active principle used in the treatment of osteoporosis and was described for the first time in European patent application EP62503. In this prior patent various preparation methods are described which generally involve the following stages:


1) protection of the 2 hydroxylic functions of 6-hydroxy-2-(4-hydroxyphenyl)benzo[b]thiophene according to the following reaction scheme

20



25 where R_5 is an alkyl, cycloalkyl or COR_6 acyl group, a SO_2R_6 sulfonyl group where R_6 is a primary or secondary C_1 - C_4 alkyl, C_1 - C_3 fluoro alkyl or C_1 - C_4 alkoxyphenyl,
2) acylation of the compound protected with 4-(2-piperidinoethoxy)benzoyl halide

according to the following synthesis scheme:



10 in which R_7 is a halogen atom,

3) deprotection or elimination of the OR_5 protective group.

As it results from the examples reported in EP62503, when the reaction is conducted using the acetyloxy group as OR_5 protective group, deprotection of this group is conducted first with sodium hydroxide in an alcoholic solution and subsequently with methanesulfonic acid. This type of hydrolysis however does not allow high purity raloxifene to be obtained, since, as indicated by example 6, the product to be purified must be passed through a chromatographic column.

15 This type of treatment, however, only enables a yellow foam to be obtained, and, to arrive at a product of solid crystalline form, a further treatment with acetone is required. The crystallized product thus obtained consisting of raloxifene methanesulfonate must be further converted into the corresponding hydrochloride for pharmaceutical use.

20

The aforesaid process, requiring product passage through a chromatographic column, is not achievable at industrial level, proof of which being that in the same prior patent, instead of the aforesaid synthesis scheme, the one preferred is that in which the OR_5 protective group is an alkoxy, specifically a methoxy group, which for unblocking requires the use of aluminium trichloride and a thioderivative and preferably methanethiol, moreover in a quantity greatly in excess of the substrate on which the deprotection must be conducted, with considerable

25

pollution problems, which evidently involves the use of considerable quantities of thioderivatives.

The processes described in EP62503 involve another inconvenience caused by the use of aluminium trichloride and, if proceeding to the scheme preferred by this prior patent, this Lewis acid must be used in substantial quantities, since it is used not only in stage (2) of acylation, but also in subsequent dealkylation. Aluminium trichloride as shown in the subsequent patent US5629425 produces a large quantity of aluminium-based by-products which are soluble in raloxifene processing solvents and are found therefore in the final product.

To overcome these problems, in the aforesaid US5629425 boron trichloride or boron tribromide is used as Lewis acid, these being decidedly more expensive catalysts than aluminium trichloride.


The need was felt to provide a process which enabled raloxifene hydrochloride to be prepared with high yields and high purity and low aluminium content without using expensive catalysts.

Summary of the invention

The applicant has surprisingly found a process capable of overcoming the drawbacks of known processes and which allows raloxifene and in particular raloxifene hydrochloride to be obtained with high purity and high yields.

This process comprises in particular the following stages:

a) demethylation of 6-methoxy-2-(4-methoxyphenyl)benzo[b]thiophene of formula (II)




(II)

in pyridine hydrochloride to obtain 6-hydroxy-2-(4-hydroxyphenyl)benzo[b]thiophene of formula (III)




b) acetylation of 6-hydroxy-2-(4-hydroxyphenyl)benzo[b]thiophene with an
 5 acetylating agent to obtain the corresponding 6-acetoxy-2-(4-acetoxyphenyl)benzo[b]thiophene of formula (IV)



(IV)


c) acylation of 6-acetoxy-2-(4-acetoxyphenyl)benzo[b]thiophene (IV) with 4-(2-piperidinoethoxy)benzoylchloride hydrochloride of formula (V)



HCl

(V)

20 with aluminium trichloride in halogenated solvent to obtain 6-acetoxy-2-(4-acetoxyphenyl)-3-[4-(2-piperidinoethoxy)benzoyl]-benzo[b]thiophene of formula (VI)




d) hydrolysis of 6-acetoxy-2-(4-acetoxyphenyl)-3-[4-(2-piperidinoethoxy)benzoyl]-benzo[b]thiophene, according to the following operative methods:

- 5 d1) treatment of 6-acetoxy-2-(4-acetoxyphenyl)-3-[4-(2-piperidinoethoxy)benzoyl]-benzo[b]thiophene with alkaline hydroxide in alcohol solvent,
 d2) acidification of the product obtained in the previous stage (d1) with a strong acid, to obtain the corresponding raloxifene salt with strong acid, characterised in that the strong acid used in stage (d2) is concentrated hydrochloric acid.
- 10 In this respect, by conducting the hydrolysis of 6-acetoxy-2-(4-acetoxyphenyl)-3-[4-(2-piperidinoethoxy)benzoyl]-benzo[b]thiophene with sodium hydroxide and subsequently treating the product obtained with hydrochloric acid in place of methanesulfonic acid, raloxifene hydrochloride precipitates in crystalline form directly with a high purity equal to 98%, thus in contrast to the analogous process
- 15 described in EP65203 conducted with methanesulfonic acid, without having to use purification processes such as passage through a chromatographic column, which are impractical from the industrial point of view. In addition the product derived from stage (d2) has a low aluminium content.

Detailed description of the invention

- 20 The 6-methoxy-2-(4-methoxyphenyl)benzo[b]thiophene of formula (II) used in stage (a) of the process of the present invention is prepared by reacting 3-methoxybenzene-thiol with α -bromo-4-methoxyacetophenone to obtain the corresponding α -(3-methoxyphenylthio)-4-methoxyacetophenone which is finally cyclized to obtain the intermediate (II) with polyphosphoric acid, as in the
- 25 following scheme.



The pyridine hydrochloride used in stage (a) is preferably prepared in situ by
5 adding concentrated hydrochloric acid to pyridine and distilling off all the water to
obtain a thick but stirrable residue. The applicant has also surprisingly found that
if the demethylation reaction or stage (a) of the process of the present invention is
conducted in the presence not only of pyridine hydrochloride but also of
tributylamine, preferably in weight ratios with respect to 6-methoxy-2-(4-
10 methoxyphenyl)benzo[b]thiophene (II) of between 0.5 and 2, it is possible to lower
the reaction temperature which in prior art is conducted at 210°C, to decidedly
lower temperatures, between 170 and 180°C.

According to a preferred embodiment of the process of the present invention, it is
not necessary to isolate the 6-hydroxy-2-(4-hydroxyphenyl)benzo[b]thiophene (III)
15 obtained in stage (a).

In stage (b) according to a preferred embodiment acetic anhydride is used as
acetylating agent and a tertiary aliphatic amine, preferably triethylamine, is used
as hydrogen ion acceptor. The solvent used in stage (a) is an aprotic polar
solvent, ethyl acetate being particularly preferred.

20 The 4-(2-piperidinoethoxy)benzoylchloride hydrochloride of formula (V) used in
stage (c) is preferably prepared in situ by a conventional type procedure by
reacting 4-(2-piperidinoethoxy)-benzoic acid hydrochloride with thionyl chloride
without isolating the reaction product. This reaction is preferably conducted in
methylene chloride in the presence of pyridine as catalyst.

25 Stage (c) is preferably conducted in methylene chloride, according to a
particularly preferred embodiment this stage being conducted in the following
manner: 6-acetoxy-2-(4-acetoxyphenyl)benzo[b]thiophene is added to 4-(2-
piperidinoethoxy)benzoylchloride hydrochloride of formula (V) prepared in situ
while still in its reaction solvent methylene chloride, the mixture thus obtained
30 being poured onto a mixture consisting of methylene chloride and aluminium

trichloride.

According to a preferred embodiment of the process of the present invention, 6-acetoxy-2-(4-acetoxyphenyl)-3-[4-(2-piperidinoethoxy)benzoyl]-benzo[b]thiophene (VI) is not isolated but is used in crude form for the subsequent hydrolysis (d).

- 5 Stage (d1) is preferably conducted using methanol as the alcoholic solvent, with excess 30% sodium hydroxide.

Stage (d2) is preferably conducted directly on the reaction mixture derived from stage (d1) to which equal weight quantities of water and ethyl acetate are added and finally 37% concentrated hydrochloric acid.

- 10 A suspension is hence obtained, which is preferably washed with equal weight quantities of water and ethyl acetate.

By the process of the present invention raloxifene hydrochloride is obtained with high purity and high yields of about 65-70% calculated on the 6-methoxy-2-(4-methoxyphenyl)benzo[b]thiophene (II).

- 15 The applicant has also found that if raloxifene hydrochloride obtained by the process of the present invention is crystallised from an alcoholic solvent, preferably methanol, possibly in the presence of small quantities of HCl, it achieves a purity of greater than 99%.

- Finally the applicant has also found that by conducting a further crystallization, again from an alcoholic solvent, preferably methanol, possibly in the presence of HCl, on the product derived from the first crystallisation, raloxifene hydrochloride can be obtained with a purity greater than 99.7%. In particular raloxifene hydrochloride obtained after the first and/or the second crystallisation contains the characteristic impurity consisting of raloxifene hydrochloride N-oxide in a quantity less than 0.05% and preferably less than 0.01%, this product also having an aluminium content less than 5 ppm.
- 20
- 25

- The product thus obtained has a particle size distribution (after gentle grinding conducted with the aim of simply homogenising the product) such that D(0.9) is $\leq 100\mu\text{m}$ and D(0.5) $\geq 40\mu\text{m}$. By further sieving a raloxifene hydrochloride is obtained with the following particle size distribution: D(0.9) between 50 and $65\mu\text{m}$ and D[4.3] $\geq 20\mu\text{m}$.
- 30

Some illustrative but non-limiting examples of the preparation process for raloxifene hydrochloride of the present invention and its relative intermediates are given.

EXAMPLE 1

5 *Preparation of 6-acetoxy-2-(4-acetoxyphenyl)benzo[b]thiophene (IV).*

24 kg of pyridine (0.303 kmol) and 28.8 kg of 37% hydrochloric acid (0.292 kmol) are fed into a reactor. The reactor is placed under vacuum and all the water is distilled off until a thick but stirrable residue is obtained.

The residue is then redissolved in 6 kg of tributylamine and 6 kg of 6-methoxy-2-
10 (4-methoxyphenyl)benzo[b]thiophene (0.022 kmol). The mixture is heated to 170-180°C and is maintained at this temperature for some hours. It is then cooled to 50-60°C and 24 kg of ethyl acetate and 60 kg of deionised water are fed into the reactor. The mixture is stirred for 15 minutes and the phases are separated. The solvent is distilled off from the organic phase under vacuum and the residue is
15 redissolved with 24 kg of ethyl acetate and 5.3 kg of triethylamine (0.052 kmol). The mixture obtained is heated to 60-65°C while being stirred and 8.9 kg of acetic anhydride (0.087 kmol) are added. The reaction mixture is stirred for 1 hour at the same temperature then is cooled to 25-30°C and 24 kg of deionised water are added. The suspension is centrifuged, washed with 6 kg of deionised water and
20 6 kg of ethyl acetate.

The product is then dried at 50-60°C and about 6.6 kg of dried product are obtained. The reaction yield is 91.1%.

EXAMPLE 2

Preparation of crude raloxifene hydrochloride.

25 PHASE A

42 kg of methylene chloride and 7.8 kg of 4-(2-piperidinoethoxy)-benzoic acid hydrochloride (0.027 kmol), 0.12 kg pyridine (0.0015 kmol) are fed into a reactor and heated under reflux and then 3.96 kg of thionyl chloride (0.033 kmol) are added. The mixture is stirred for 1 hour then about 20 litres of methylene chloride
30 are distilled off. The mixture is cooled to 20-30°C and 6 kg of 6-acetoxy-2-(4-acetoxyphenyl)benzo[b]thiophene (IV) (0.018 kmol) are added.

The mixture is stirred until is completely homogenised.

PHASE B

36 kg of methylene chloride and 16.8 kg of aluminium trichloride (0.126 kmol) are fed into a reactor.

- 5 While stirring, the chloromethylene suspension, comprised of phase A prepared as described above, is added at 15-30°C. The mixture is stirred for 1 hour then the entire reaction mixture is poured into a reactor containing 60 kg of ice.

The mixture is stirred at 15-30°C then the suspension is centrifuged, washing with 3 kg of methylene chloride and 3 kg of deionised water.

- 10 The centrifuged mother liquors, containing the product, are fed into a reactor and the phases are separated. The organic phase is distilled off until obtaining an oily residue and 15 kg of methyl alcohol are added, stirred at 20-40°C and, maintaining the same temperature, 9.1 kg of 30% sodium hydroxide (0.068 kmol) are poured in. The mixture is stirred for 1 hour and 30 kg of deionised water and
15 30 kg of ethyl acetate are added.

- At the same temperature 7.2 kg of 37% hydrochloric acid (0.073 kmol) are then added. The suspension is centrifuged, washing with 6 kg of ethyl acetate and 6 kg of deionised water. At the end 6.6 kg of dried product with HPLC purity > 98% and low aluminium content are obtained. The reaction yield calculated on the 6-
20 acetoxy-2-(4-acetoxyphenyl)benzo[b]thiophene (IV) is equal to a yield of 70.4%.

EXAMPLE 3

Crystallisation of crude raloxifene hydrochloride (1st crystallisation of crude raloxifene hydrochloride)

- 6 kg of deionised water, 6 kg of crude raloxifene hydrochloride prepared as
25 described in example 2 and 107 kg of methyl alcohol are fed into a reactor. The reaction mixture is heated until a complete solution is obtained then 0.25 kg of decolourising carbon are added. It is stirred for 15 minutes and then the suspension is filtered. While maintaining the solution stirred, 67 kg of methyl alcohol are distilled off. The residue is cooled and 0.1 kg of 37% hydrochloric
30 acid are added. The pH, which must not exceed 2, is checked and the reaction mixture is then stirred for 2 hours at 20-40°C. The suspension is centrifuged,

washing with 6 kg of methyl alcohol. 4.5 kg of dried product are obtained with HPLC purity of >99% and a yield of 75%.

EXAMPLE 4

Crystallisation of crystalline raloxifene (2nd crystallisation).

5 0.9 kg of deionised water, 81 kg of methanol and the entire amount of crystallised product as described in example 3 are fed into a reactor. While maintaining the reaction mixture under stirring it is heated under reflux and 36 kg of methyl alcohol are distilled off. It is then cooled to 20-40°C and 0.08 kg of 37 % hydrochloric acid are added. The suspension is centrifuged, washing with 4 kg of
10 methyl alcohol. The product is dried at 70°C. 4 kg of raloxifene hydrochloride are obtained with HPLC purity > 99.8%, reaction yield 89%, in particular the raloxifene hydrochloride N-oxide content is less than 0.01% and aluminium content is less than 5ppm. In particular the raloxifene hydrochloride obtained after crystallisation
15 in a quantity less than 0.05% and preferably less than 0.01%. The product thus obtained has a particle size distribution (after gentle grinding conducted with the aim of simply homogenising the product) such that D(0.9) is $\leq 100\mu\text{m}$ and D (0.5) $\geq 40\mu\text{m}$.


By further sieving a raloxifene hydrochloride is obtained with the following particle
20 size distribution: D(0.9) between 50 and 65 μm and D[4.3] $\geq 20\mu\text{m}$.

11

CLAIMS


1. Process for preparing raloxifene hydrochloride of formula (I)

5




with a purity higher than 98% comprising the following stages:

- 10 a) demethylation of 6-methoxy-2-(4-methoxyphenyl)benzo[b]thiophene of formula (II)



15

in pyridine hydrochloride to obtain 6-hydroxy-2-(4-hydroxyphenyl)benzo[b]thiophene of formula (III)




20

b) acetylation of 6-hydroxy-2-(4-hydroxyphenyl)benzo[b]thiophene with an acetylating agent to obtain the corresponding 6-acetoxy-2-(4-acetoxyphenyl)benzo[b]thiophene of formula (IV)


25

12



(IV)


c) acylation of 6-acetoxy-2-(4-acetoxyphenyl)benzo[b]thiophene (IV) with 4-(2-piperidinoethoxy)benzoylchloride hydrochloride of formula (V)



HCl

(V)

with aluminium chloride in halogenated solvent to obtain 6-acetoxy-2-(4-acetoxyphenyl)-3-[4-(2-piperidinoethoxy)benzoyl]-benzo[b]thiophene of formula (VI)



(VI)

d) hydrolysis of 6-acetoxy-2-(4-acetoxyphenyl)-3-[4-(2-piperidinoethoxy)benzoyl]-benzo[b]thiophene, according to the following operative modalities:

- d1) treatment of 6-acetoxy-2-(4-acetoxyphenyl)-3-[4-(2-piperidinoethoxy)benzoyl]-benzo[b]thiophene with alkaline hydroxide in alcohol solvent,
- d2) acidification of the product obtained in the preceding stage (d1) with a strong acid, to obtain the corresponding raloxifene salt with the strong acid,
- 5 characterised in that the strong acid used in stage (d2) is concentrated hydrochloric acid.
2. Process as claimed in claim 1, characterised in that the pyridine hydrochloride used in stage (a) is prepared in situ by adding concentrated hydrochloric acid to pyridine and distilling off all the water to obtain a thick but stirrable residue.
- 10 3. Process as claimed in claim 1 or 2, characterised in that the demethylation reaction or stage (a) of the process of the present invention is also conducted in the presence or tributylamine.
4. Process as claimed in claim 3, characterised in that tributylamine is used preferably in weight ratios with respect to 6-methoxy-2-(4-methoxyphenyl)benzo[b]thiophene (II) of between 0.5 and 2.
- 15 5. Process as claimed in claim 4, characterised in that stage (a) is conducted at a temperature between 170 and 180°C.
6. Process as claimed in any one of claims 1-5, characterised in that acetic anhydride is used as acetylating agent in the presence of triethylamine in ethyl acetate.
- 20 7. Process as claimed in any one of claims 1-6, characterised in that the 4-(2-piperidinoethoxy)benzoylchloride hydrochloride of formula (V) used in stage (c) is prepared in situ, by reacting 4-(2-piperidinoethoxy)benzoic acid hydrochloride with thionyl chloride in methylene chloride in the presence of pyridine, without isolating the reaction product.
- 25 8. Process as claimed in any one of claims 1-7, characterised in that stage (c) is conducted in methylene chloride.
9. Process as claimed in claim 8, characterised in that stage (c) is conducted according to the following operative modalities: 6-acetoxy-2-(4-acetoxyphenyl)benzo[b]thiophene (IV) is added to non-isolated 4-(2-piperidinoethoxy)benzoylchloride hydrochloride of formula (V) and prepared in
- 30

situ as in claim 7 and the aforesaid mixture is poured onto a mixture consisting of methylene chloride and aluminium trichloride.

10. Process as claimed in any one of claims 1-9, characterised in that the 6-acetoxy-2-(4-acetoxyphenyl)benzo[b]thiophene (IV) is not isolated, but is used in
5 the crude state in the subsequent reaction (d).

11. Process as claimed in any one of claims 1-10, characterised in that stage (d1) is conducted using methanol as alcohol solvent and excess 30% sodium hydroxide.

12. Process as claimed in any one of claims 1-11, characterised in that stage (d2)
10 is conducted directly on the reaction mixture derived from stage (d1) to which are added equal weight quantities of water and ethyl acetate and finally 37% concentrated hydrochloric acid.

13. Process as claimed in claim 1-12, characterised in that the suspension obtained in stage (d2) is washed with equal weight quantities of water and ethyl
15 acetate.

14. Process as claimed in any one of claims 1-13, characterised in that raloxifene hydrochloride has an HPLC purity >98%.

15. Process as claimed in any one of claims 1-14, characterised in that raloxifene hydrochloride derived from stage (d2) is crystallised from an alcoholic solvent.

20 16. Process as claimed in claim 15, characterised in that said solvent is methanol possibly in the presence of HCl.

17. Process as claimed in any one of claims 15 and 16, characterised in that raloxifene hydrochloride is obtained with a purity greater than 99%.

18. Process as claimed in any one of claims 15 and 16, characterised in that a
25 further crystallisation from raloxifene hydrochloride from alcohol solvent is conducted.

19. Process as claimed in claim 18, characterised in that said crystallisation is conducted in methanol possibly in the presence of HCl.

20. Raloxifene hydrochloride with a purity greater than 99.7%.

30 21. Raloxifene hydrochloride as claimed in any one of claims 17 – 20, characterised in that it contains aluminium in a quantity less than 5 ppm.

22. Raloxifene hydrochloride as claimed in any one of claims 17-21 characterised in that it contains raloxifene hydrochloride N-oxide in a quantity less than 0.05%.

23. Raloxifene hydrochloride as claimed in claim 22, characterised in that said impurity is contained in a quantity less than 0.01%.

5 24. Raloxifene hydrochloride as claimed in any one of claims 20-23, characterised by having a $D(0.9) \leq 100\mu\text{m}$ and a $D(0.5) \geq 40\mu\text{m}$.

25. Raloxifene hydrochloride as claimed in claim 24, characterised, after a further sieving, by having a $D(0.9)$ between 50 and 65 μm and a $D[4.3] \geq 20\mu\text{m}$.

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/EP2004/051263

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D333/56

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the International search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, BEILSTEIN Data, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 0 062 503 A (LILLY CO ELI) 13 October 1982 (1982-10-13) cited in the application preparations 1,5 page 41, line 13 - line 16; examples 2,6,17	1-19
X	example 20	20-25
X	US 4 418 068 A (JONES CHARLES D) 29 November 1983 (1983-11-29) examples 15,18	20-25

☐ Further documents are listed in the continuation of box C.

Patent family members are listed in annex.

* Special categories of cited documents:

A document defining the general state of the art which is not considered to be of particular relevance

E earlier document but published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

& document member of the same patent family

Date of the actual completion of the international search

15 October 2004

Date of mailing of the international search report

25/10/2004

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

De Jong, B

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

EP2004/051263

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
EP 0062503	A	13-10-1982	AT 17243 T	15-01-1986
			AT 14429 T	15-08-1985
			AU 555658 B2	02-10-1986
			AU 8226582 A	07-10-1982
			BG 37378 A3	15-05-1985
			CS 227348 B2	16-04-1984
			DD 201794 A5	10-08-1983
			DD 201793 A5	10-08-1983
			DE 3264853 D1	29-08-1985
			DE 3268227 D1	13-02-1986
			DK 151282 A	04-10-1982
			DK 151382 A	04-10-1982
			EG 15610 A	30-06-1986
			EP 0062503 A1	13-10-1982
			EP 0062504 A1	13-10-1982
			EP 0062505 A1	13-10-1982
			ES 8304116 A1	16-05-1983
			ES 8307240 A1	16-10-1983
			FI 821160 A	04-10-1982
			FI 821161 A	04-10-1982
			GB 2097392 A ,B	03-11-1982
			GB 2097788 A ,B	10-11-1982
			GB 2096608 A ,B	20-10-1982
			GR 76468 A1	10-08-1984
			GR 78364 A1	26-09-1984
			HU 191084 B	28-01-1987
			HU 191353 B	27-02-1987
			IE 53187 B1	17-08-1988
			IL 65378 A	28-02-1986
			JP 57181081 A	08-11-1982
			KR 8501881 B1	28-12-1985
			KR 8501882 B1	28-12-1985
			MX 9203117 A1	01-07-1992
			NZ 200204 A	31-05-1985
			PH 18628 A	23-08-1985
			PL 235751 A1	22-11-1982
			PL 235752 A1	22-11-1982
			PT 74692 A ,B	01-05-1982
			PT 74693 A ,B	01-05-1982
			RO 84584 A1	17-07-1984
			SU 1155157 A3	07-05-1985
			US 4418068 A	29-11-1983
			ZA 8202247 A	30-11-1983
US 4418068	A	29-11-1983	MX 9203117 A1	01-07-1992
			AT 17243 T	15-01-1986
			AT 14429 T	15-08-1985
			AU 555658 B2	02-10-1986
			AU 8226582 A	07-10-1982
			BG 37378 A3	15-05-1985
			CS 227348 B2	16-04-1984
			DD 201794 A5	10-08-1983
			DD 201793 A5	10-08-1983
			DE 3264853 D1	29-08-1985
			DE 3268227 D1	13-02-1986
			DK 151282 A	04-10-1982
			DK 151382 A	04-10-1982
			EG 15610 A	30-06-1986

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP2004/051263

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 4418068 A		EP 0062503 A1	13-10-1982
		EP 0062504 A1	13-10-1982
		EP 0062505 A1	13-10-1982
		ES 8304116 A1	16-05-1983
		ES 8307240 A1	16-10-1983
		FI 821160 A	04-10-1982
		FI 821161 A	04-10-1982
		GB 2097392 A ,B	03-11-1982
		GB 2097788 A ,B	10-11-1982
		GB 2096608 A ,B	20-10-1982
		GR 76468 A1	10-08-1984
		GR 78364 A1	26-09-1984
		HU 191084 B	28-01-1987
		HU 191353 B	27-02-1987
		IE 53187 B1	17-08-1988
		IL 65378 A	28-02-1986
		JP 57181081 A	08-11-1982
		KR 8501881 B1	28-12-1985
		KR 8501882 B1	28-12-1985
		NZ 200204 A	31-05-1985
		PH 18628 A	23-08-1985
		PL 235751 A1	22-11-1982
		PL 235752 A1	22-11-1982
		PT 74692 A ,B	01-05-1982
		PT 74693 A ,B	01-05-1982
		RO 84584 A1	17-07-1984
		SU 1155157 A3	07-05-1985
		ZA 8202247 A	30-11-1983